



Washington Township Medical Foundation

Part of Washington Hospital Healthcare System

Washington Women's Health

2299 Mowry Avenue, Suite 3C

Fremont, California 94538

Phone: 510-248-1470

Fax: 510-796-5198

Stacey Barrie, MD
Siobhan Calhoun, MD
Jing Dai, MD
Elizabeth Kurkjian, MD
Victoria Leiphart, MD
Jelriza Mansouri, MD
Stacie Macdonald, MD
Sudeepthi Prasad, MD
Jenny (Hanh Thi) Tran, DO

Diplomates of the American Board
of Obstetrics and Gynecology

Post-Operative Instructions

Patient name:

Operation(s):

Recovery following minor gynecologic surgery generally occurs rapidly, with minimal discomfort. You should rest for the remainder of today. Then, gradually increase activities as tolerated. The following instructions will guide you through your recovery.

- You may return to work in _____ day(s).
- If you exercise regularly, gradually resume this in one week unless instructed otherwise.
- Abstain from intercourse, douching, or placing anything in your vagina for _____ days.

Remove Band-Aids tomorrow morning. Showers are permitted thereafter. Replace Band-Aids only to absorb discharge from incisions. Leave underlying steri-strips in place until they fall off or a maximum of one week.

Please be aware of the following potential problems:

Pain:

Take _____, _____ tablets every _____ hours as needed. You may also use Motrin® (Ibuprofen) 600mg every 6 hours, which is available without prescription. If, despite medication your pain grows stronger and becomes unbearable, call your doctor. Do not drive or operate dangerous machinery until off pain medication for greater than 24 hours.

Bleeding:

This will vary depending on the operation. Vaginal bleeding should not exceed a heavy menstrual period. If bleeding consistently saturates a pad every hour, call your doctor.



Washington Township Medical Foundation

Part of Washington Hospital Healthcare System

Washington Women's Health

2299 Mowry Avenue, Suite 3C
Fremont, California 94538
Phone: 510-248-1470
Fax: 510-796-5198

Stacey Barrie, MD
Siobhan Calhoun, MD
Jing Dai, MD
Elizabeth Kurkjian, MD
Victoria Leiphart, MD
Jelriza Mansouri, MD
Stacie Macdonald, MD
Sudeepthi Prasad, MD
Jenny (Hanh Thi) Tran, DO

Diplomates of the American Board
of Obstetrics and Gynecology

Infection:

If you feel warm or chilled, check your temperature. If you have a persistent fever greater than 101.5 F or 38.5 C for more than 6 hours, call your doctor. In addition, if you have a skin incision which becomes increasingly red or leaks pus, call your doctor.

Bladder:

You should be able to empty your bladder every 6 hours. Please call your doctor if you are unable to void.

Bowels:

Bowel movements may be delayed for a few days. You may use a laxative, like Milk-of-Magnesia according to package instructions, if constipation continues past the 3rd day after surgery.

Special Instructions:

If you have not already done so, please schedule a post-operative appointment with our office for approximately 2 weeks from today.

(Physician)

(Date)